

READ THE LABEL BEFORE USE. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. SHAKE THE BOTTLE WELL BEFORE USE



Plant Growth Regulator

ETHEPHON 480 SL

Reg. No. L 4776 Act 36 of 1947
N-AR 0994 / W1301438

8: 27/08/2024 – Oct2024

A soluble concentrate plant growth regulator
for use on crops as indicated.

ACTIVE INGREDIENT

ethephon 480 g/l



DANGER

Hazard Statements:

May be harmful if swallowed.
Harmful if contact with skin.
Causes severe skin burns and eye damage.
Toxic to aquatic life with long lasting effects.

Precautionary Statements:

Do not breathe dusts or mists.
Wash hands and face thoroughly after handling. Avoid release into the environment.
Wear impervious rubber gloves and boots, protective clothing, and chemical safety goggles.



villa 

UN Number: 2922

Registration holder: VILLA CROP PROTECTION (PTY) LTD.
Co. Reg. No. 1992/002474/07
65 Botes Road, Glen Marais, Kempton Park, 1619
Tel. (011) 396 2233
Website: www.villacrop.co.za

IN CASE OF POISONING / 24 HR EMERGENCY NUMBERS:
Griffon Poison Information Centre (National): +27 82 446 8946
Poison Information Helpline (National): +27 861 555 777
24 Hr Transport / Spill emergency no: (Hazcall24) +27 86 044 4411
(Client: Villa Crop Protection)

REFER TO DETAILS AS PRINTED ON CONTAINER / BAG

DIRECTIONS FOR USE ENCLOSED

Batch Number:
Date of Manufacture:

ETHEPHON 480 SL

Reg. No. L 4776 Act No. 36 of 1947

N-AR 0994 / W1301438

ACTIVE INGREDIENT:

ethephon 480 g/l

Registration holder:

VILLA CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1992/002474/07

65 Botes Road, KEMPTON PARK, 1619

Tel. (011) 396 2233

WARNINGS**Hazard Statements:**

May be harmful if swallowed.
Harmful in contact with skin.
Causes severe skin burns and eye damage.
Toxic to aquatic life with long lasting effects

- **IMPORTANT - EXPORT CROPS**

IN THE CASE OF CROPS INTENDED FOR EXPORT, CONFIRM WITHHOLDING PERIODS WITH THE RELEVANT EXPORTING ORGANISATION. IF THIS PRODUCT IS USED ACCORDING TO THE RECOMMENDATIONS ON THIS LABEL, THE LOCAL MRL VALUES WILL NOT BE EXCEEDED. HOWEVER, VILLA CROP PROTECTION (PTY) LTD. CANNOT ACCEPT RESPONSIBILITY FOR EXPORTED CROPS EXCEEDING THE IMPORT TOLERANCES OF OTHER COUNTRIES.

- Handle with care.
- Product is corrosive.
- Store away from food, feeds, seed and other agricultural chemicals.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Aerial application:

Notify all inhabitants in the immediate area to be sprayed and issue the necessary warnings. Do not spray over or allow drift to contaminate water or adjacent areas.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS**Precautionary statements:**

Do not breathe dusts or mists.
Wash hands and face thoroughly after handling.
Avoid release to the environment.
Wear impervious rubber gloves and boots, protective clothing and chemical safety goggles.
IF SWALLOWED: Get medical help.
IF SWALLOWED: Rinse mouth. Do NOT Induce vomiting.
IF ON SKIN: Take off immediately all contaminated clothing. Immediately rinse with water for several minutes.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
Get medical help.
Collect spillage.
Store locked up.
Dispose of content/container to suitable landfill in accordance with local regulations

- Wash contaminated clothing after use.
- Wash with soap and water after use or after accidental skin contact.
- Do not eat, drink or smoke while mixing, applying or before washing hands and face or change of clothing.
- Prevent spray drift and/or contamination onto susceptible crops, grazing, rivers, dams and any other areas not under treatment.
- Clean application equipment after use. Do not dispose of wash water where it can contaminate other crops, grazing, boreholes, rivers or dams.
- **TRIPLE RINSE THE EMPTY CONTAINER AS FOLLOWS:** Invert the empty container over the spray or mixing tank and drain for at least 30 seconds after the flow has slowed down to dripping. Thereafter rinse the empty container three times in succession with one quarter of the container volume fresh water and decant the rinsate into the spray or mixing tank. Puncture the triple rinsed container and dispose of via an approved collector or recycler www.croplife.co.za. Do not bury, burn, or donate the container to any other parties that may use it as a container for food or beverages.
- Prevent contamination of food, feeds, drinking water and eating utensils.

FIRST AID TREATMENT

- Remove the victim from the area of exposure. Wash off remaining material with plenty of water. In the event of any complaints or symptoms, avoid further exposure and consult a doctor.
- **Inhalation:** Remove person from contaminated area to fresh air and assist breathing as needed. **Seek medical attention if you feel unwell after inhalation)**
- **Skin:** Remove contaminated clothing and shoes. Gently wipe off excess chemical. Wash skin gently and thoroughly with water and non-abrasive soap. **Immediately seek medical attention if irritation persists.**
- **Eyes:** Flush eyes with clean water for at least 15 – 20 minutes. Lift eyelids to facilitate irrigation. If present, remove contact lenses after 5 minutes and continue rinsing. **Seek medical attention immediately.**
- **Ingestion:** Seek medical attention or call a poison control centre for treatment advice. May cause gastrointestinal irritation or ulceration. Due to the solvent, this product is classified as an aspiration hazard. Mild to severe pulmonary injury could be caused due to the aspiration of small amounts of product during vomiting or ingestion. Do not induce vomiting unless instructed to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. If the person is alert, rinse mouth thoroughly with water. **Seek medical attention immediately.**

Relevant hazardous components	
Ethephon TC	480 g ae/l

USE RESTRICTIONS

- Spray deposits of **ETHEPHON 480 SL** plant growth regulator, if left on acrylic, plastic type aircraft windshields for more than one hour, may react with this material.
- An opaque film or spots may form on the surface, which cannot be removed by washing or with any other cleaner.
- Windshields should be rinsed with water after each tank load of **ETHEPHON 480 SL** has been sprayed.
- The aircraft should be rinsed thoroughly each day after spraying **ETHEPHON 480 SL**.

DIRECTIONS FOR USE: Use only as directed.

ETHEPHON 480 SL must only be applied with a calibrated sprayer that is in good working order.

Mixing instructions:

- Fill the spray tank with half of the required amount of water.
- Add the required amount of **ETHEPHON 480 SL** while agitating.
- Add the remaining amount of water.
- Ensure thorough agitation of the mixture in the tank during mixing and spraying.
- Prepare only as much spray solution as can be used on one day of spraying.
- Do not allow the spray solution to stand overnight.
- Thoroughly flush out the spraying equipment at the end of the spraying operation.

Post Spray Equipment Cleaning

It is essential to rinse the spray tank and all hoses with a 0.1 % solution of either calcium chloride or ammonium hydroxide or the recommended rate of **Protank® liquid cleaner** according to the product label. Allow this solution to stand in the spray equipment for 15 minutes; empty the spray equipment; repeat the rinsing with a 0.1 % solution of calcium chloride or ammonium hydroxide for 15 minutes; rinse all equipment

with water. Nozzles and fitters must be cleaned individually. Rinse water and solution should be drained in a cesspool or drain where it will not contaminate any water source.

Compatibility:

ETHEPHON 480 SL is compatible with **Villa 51** and **Masterlock®**. The full range of physical compatibility of **ETHEPHON 480 SL** with other pesticides and fertilizers is not known. Therefore, before preparing commercial tank mixes, add a small amount of **ETHEPHON 480 SL** to a proportional volume of water and then the pesticide. Do not use mixtures that curdle, precipitate or grease.

Early detection of latent Citrus black spot (CBS):

- Samples must be collected eight (8) weeks prior to harvest from sensitive indicator trees, within each orchard (sensitive indicator trees are trees in a poor condition and trees at rows ends where there is a concentration of risk of deficient spray application).
- The sample size should be 100 fruit per orchard (up to 5-hectare orchards).
- If an orchard exceeds 5 hectares, the sample size should be increased by 20 fruit per hectare greater than 5 hectares.
- Half of the sample (50 fruit) should be collected from row end trees, evenly spread across both ends of the orchard.
- The remaining sample (50 fruit) should be collected from trees in poor condition. If there are less than 10 trees in poor condition, the shortfall should be corrected by collecting fruit from trees evenly spaced in a diagonal transect across the orchard.
- From each sample tree, collect the fruit from the lower outside portion of the tree canopy on the sunny side of the tree.
- **All the treated fruit must be disposed of and none of them may be consumed.**

Aerial Application:

Aerial application of **ETHEPHON 480 SL** may only be done by a registered aerial application operator using a correctly calibrated, registered aircraft according to the instructions of SANS Code 10118 (Aerial Application of Agricultural Pesticides). Ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria be met:

- Volume: A minimum spray mixture volume of 30 litres per hectare is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aerially at a lower volume rate than recommended above.
- Droplet coverage: 25 to 35 droplets per cm² must be recovered at the target area.
- Droplet size: A droplet spectrum with a VMD of 280 to 300 micron is recommended. Limit the production of fine droplets less than 150 micron (high drift and evaporation potential) to a minimum.
- Flying height: Maintain the height of the spray boom at 3 to 4 metres above the target. Do not spray when aircraft dives, is in a climb or when banking.
- Use suitable atomising equipment that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product. The spraying system must produce a droplet spectrum with the lowest possible Relative Span.
- Position all the atomisers within the inner 60 to 75 % of the wingspan to prevent droplets from entering the wingtip vortices.
- The difference in temperature between the wet and dry bulb thermometers, of a whirling hygrometer, should not exceed 8 °C.
- Stop spraying if the wind speed exceeds 15 km per hour.
- Stop spraying under turbulent, unstable and dry conditions during the heat of the day.
- Spraying under temperature inversion conditions (spraying in or above the inversion layer) and/or high humidity conditions (relative humidity 80 % and above) may lead to the following:
 - reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage),
 - damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.
- Ensure that the aerial spray operator knows exactly which fields to spray.

Obtain an assurance from the aerial spray operator that the above requirements will be met and that relevant data will be compiled in a logbook and kept for future reference.

APPLICATION RATES

Crop / Use	Dosage Rate	Remarks
Citrus Early detection of latent Citrus black spot	8 ml / 1 ℓ water	Prepare an ETHEPHON 480 SL suspension in a suitable container. Dip fruit into the suspension for five (5) minutes. Ensure all fruit are submerged. After removing the fruit from dipping, let them air-dry overnight. Store the fruit at approximately +/- 25°C under dry conditions for fourteen (14) days. Destroy fruit after inspection.
Sugarcane Increase sucrose content	1.5 ℓ / 30 ℓ water / ha	Ensure thorough coverage by means of aerial application. To obtain maximum benefit, apply only to actively growing cane 6 to 12 weeks before harvest.
Pineapples Colouring of fruit	2.0 to 4.0 ℓ / 500 to 1000 ℓ water / ha	ETHEPHON 480 SL may affect sugar concentration unless applied at or just after natural colour change. For uniform colouring, ensure a thorough coverage of fruit with the spray solution. For maximum benefit direct spray on fruit only.
Induce flowering: Cayenne's	1.0 ℓ PLUS 50 ℓ U.A.N. (32) / 2500 ℓ water / ha	Apply to plants when they are ready to be forced to flower.
Induce flowering: Queens	125 ml PLUS 1.5 ℓ U.A.N. (32)	Use 20 ml of mixture per individual plant or sucker. Apply to individual plants or suckers when they are ready to be forced to flower.
Cherries (Sour and Yellow) Uniform ripening and easier harvest	50 ml / 100 ℓ water	NOT SUITABLE FOR FRESH MARKET CHERRIES. Apply as a full cover spray approximately 10 days prior to anticipated harvest.
Wheat Reducing the incidence of lodging in susceptible varieties grown under conditions conducive to lodging (SST33, 44, 66 & 107, Elize, Inia & Palmiet)	1.0 to 1.25 ℓ / ha	Apply between stages 15 (matured flag leaf visible) and 16 (awns of the ear visible), according to the growth stages of the ARC-Small Grain Institute in Bethlehem. Ground Application: Ensure the spray boom is approximately 50 cm above the crop. Apply as a full cover spray in 250 to 450 litres water per hectare. Aerial application: Refer to " Aerial Application " instructions above.
NOTES		
<ul style="list-style-type: none"> • Ensure that insect, weed and disease control in the crop to be treated, are of a high standard. • ETHEPHON 480 SL only becomes effective 7 to 10 days after application. • The efficacy of the product may be impaired due to abnormal environmental conditions. Do not apply to wheat crops under drought stress or if rain is imminent. • Do not commence irrigation within 6 hours after application. • Allow 7 days between last application and harvest or grazing of wheat. 		
Apples Improved colour development (Starking types) and Enhancement of maturity (Starking types & Granny Smith)	75 to 85 ml / 100 ℓ water	Only for the local market. Application as a full cover spray should be performed 2 to 3 weeks before anticipated harvest. Use the lower dosage rate if high temperature conditions occur whilst spraying.

Crop / Use	Dosage Rate	Remarks
Cotton To accelerate opening of mature, unopened cotton bolls	2.0 to 3.0 ℓ / ha	Apply as full cover spray when 60 % of the bolls are open or mature. A boll is mature when it is very hard and won't dent when squeezed, is difficult to slice with a sharp knife and the seed coat is light brown in colour. Ensure bolls are well covered by the spray. Ground application: Apply in 150 to 350 litres water per hectare. Aerial application: Apply in 40 to 50 litres water per hectare. Use the lower rate only on wide spaced cotton.
NOTES <ul style="list-style-type: none"> Do not plant a follow-up crop within 30 days after treatment. Do not apply if rain is expected within 6 to 8 hours. Do not apply on cotton planted after 15 November, or before 150 days after planting or at 70 % boll opening. Some premature drop of small, immature bolls may be associated with the treatment. ETHEPHON 480 SL may stimulate the opening of bolls, which are not mature. This could lead to lower quality seed and lint, which can be minimized by closely monitoring the crop and applying ETHEPHON 480 SL when there are sufficient mature unopened bolls to produce the desired yield. 		
Peaches Enhancement of maturity Early cultivars	20 ml / 100 ℓ water	Only for Culemborg, Rhodes, Van Riebeeck and Earlibelle (the latter only in Gauteng) cultivars. Application as a full cover spray is performed 2 to 3 weeks before anticipated harvest. The addition of a registered wetting agent is recommended.
Plums Advanced and more uniform ripening	20 ml / 100 ℓ water PLUS registered wetting agent	Only for Cultivars Santa Rosa, Gaviota and Golden King cultivars. Apply as a full cover spray 2 weeks before anticipated harvest to commencement of harvest.
Grapes Colour improvement of Barlinka cultivar	50 ml / 100 ℓ water PLUS 100 ml / 100 ℓ water Masterlock®	Only for Barlinka cultivar. Apply at 5 % colouring of berries. Harvest at the correct degree of maturity, according to the sugar/acid content. Applications with ETHEPHON 480 SL can be applied with Masterlock® to ensure adequate coverage of the fruit.
Colour improvement of Flame and Crimson Seedless cultivars	40 ml / 100 ℓ water OR 40 ml / 100 ℓ water PLUS 50 to 100 g / 100 ℓ water S-Absciscic Acid PLUS 100 ml / 100 ℓ water Masterlock®	Flame and Crimson Seedless cultivars. Apply at veraison and follow up 7 days later. Harvest at the correct degree of maturity, according to the sugar/acid content. Climate conditions can play a significant role in colour development and the breakdown of the active ingredient Ethephon . Make sure you understand the conditions of the vineyard before it is treated. Ensure that the MRL levels is lower than the intended target value before picking commences. Applications with ETHEPHON 480 SL can be applied with Masterlock® to ensure adequate coverage of the fruit.

Crop / Use	Dosage Rate	Remarks
<p><u>Litchis</u></p> <p>Control of late autumn vegetative flush (End March until beginning May)</p> <p>(Cultivars Mauritius, McLean's Red & Early Delight only)</p>	<p><u>Dry autumn:</u> 104 ml / 100 l water (500 ppm active ingredient)</p> <p><u>Wet autumn:</u> 208 ml / 100 l water (1000 ppm active ingredient)</p> <p>PLUS</p> <p>100 ml / 100 l water Masterlock®</p>	<p>The time for treatment depends on various factors (such as the production area, season, and the onset of the cold period), but is usually between end of March to beginning of May.</p> <p>Apply only to young as spot-spray treatment to soft vegetative flush just after unfolding of the foliage. Apply where new vegetative flush appears during late autumn.</p> <p>ETHEPHON 480 SL will not be effective when applied later to extended vegetative growth. Ensure good coverage of the young flush foliage. Applications with ETHEPHON 480 SL can be applied with Masterlock® to ensure adequate coverage of the young flush foliage.</p> <p>Use the lower rate under dry weather conditions and the higher rate under wet weather conditions.</p> <p>Do not apply to sick or stressed trees.</p> <p>Do not apply together with any other products that is not registered.</p>
<p><u>Macadamias</u> (Cultivar Beaumont only)</p> <p>To promote abscission of nuts</p>	<p>208 ml / 100 l water</p> <p>PLUS</p> <p>0.1 % Villa 51</p> <p>PLUS</p> <p>100 ml / 100 l water Masterlock®</p>	<p>Also refer to “NOTES” below.</p> <p>Apply from March to July on trees when the nuts are physiologically mature.</p> <p><u>Mist blowers (e.g., Cima, Jacto Arbus, Nobili):</u> Apply 4 to 5 litres ETHEPHON 480 SL per hectare in 2000 to 2500 litres water per hectare.</p> <p><u>Handgun/Lance:</u> Apply 4.5 to 5.2 litres ETHEPHON 480 SL per hectare in 3000 to 3500 litres water per hectare. (Approximately 150 ml per 100 litres of water).</p> <p>Ensure thorough wetting of all nuts.</p> <p>Applications with ETHEPHON 480 SL can be applied with Masterlock® to ensure adequate coverage of the fruit.</p>
<p><u>NOTES</u></p> <ul style="list-style-type: none"> • It is important to confirm nut maturity by means of an “in-field” and “Home/Lab test”. <ul style="list-style-type: none"> - <u>In field test:</u> Check the colour of the inside of husk which should be brown to dark brown. - <u>Home/Lab test:</u> Take a sample of 100 nuts per cultivar, dehusk immediately, dry overnight in an oven/warming drawer at 90 °C (dry for longer if necessary), hand crack the next morning and float the kernels in a jug of tap water. Crop is mature if >95 % of kernels float. Note that hybrids have a lower oil percentage than Integs therefore 95 % floating will not always be reached. <p>Apply approximately 2.5 to 4.0 litres spray mixture per tree for 5 to 6-year-old trees and 6 to 8 litres per tree for mature (8 to 10-year-old trees).</p>		

Crop / Use	Dosage Rate	Remarks
Irrigated Maize Reduction of Plant (Stand) Height Limpopo & Northwest Province only	600 ml / ha	Apply as a foliar spray at V6 stage (6th leaves collar needs to be visible – BBCH 16). Apply in 300 litres water per hectare as an overall application. Ensure thorough coverage and wetting of the crop.
NOTES <ul style="list-style-type: none"> • The efficacy of the product may be impaired due to abnormal environmental conditions. Do not apply to maize crops under drought stress or if rain is imminent. • Ensure that insect, weed and disease control in the crop to be treated, are of a high standard. • Do not commence irrigation within six (6) hours after application. • Reduction in plant height is to enhance the general management practices (i.e., pesticide applications, facilitate and accelerate harvesting etc.) associated with the production of the crop. 		

The following product mentioned in this label may be replaced with an equivalent product:

- **VILLA 51** (L 8050 / W 130454 / N-AR 1090) = **WEN 51** (L 8315)

VILLA 51 and **WEN 51** are registered products of
VILLA CROP PROTECTION (PTY) LTD.

MASTERLOCK® and **PROTANK® LIQUID CLEANER** are registered trademarks of
WINFIELD SOLUTIONS REGISTRATION HOLDINGS (PTY) LTD.